

Kaput[®]

COMBO BAIT[®] PELLETS

RODENT CONTROL PELLETS

PLACE PACKS

NET WT. 3.125 LB (1.42 kg) (25 PACKS x 2 oz.)

ACTIVE INGREDIENTS:

Imidacloprid (CAS Number 138261-41-3).....0.020%
Warfarin (CAS Number 81-81-2).....0.025%

OTHER INGREDIENTS99.955%

TOTAL100.000%

Manufactured by:
Scimetrics

LTD. CORP.
P.O. Box 1045
Wellington, CO 80549

New Solutions to Old Problems

Kills Norway Rats, Roof Rats, House Mice and Voles.
First dead rats, mice and voles will appear
4 to 5 days after feeding begins.

A Bait Containing a Rodenticide to Kill Norway Rats, Roof
Rats, House Mice and Voles and an Insecticide to Aid in the
Reduction of Flea Populations on Listed Rodent Hosts



EPA Est. 72500-CO-1
EPA Reg. No. 72500-13

Keep Out of Reach of Children

CAUTION

Harmful If Swallowed.

See back panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

If swallowed, call a poison control center, doctor or 1-800-858-7378 immediately for treatment advice. Have person spit a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

NOTICE TO PHYSICIAN OR VETERINARIAN

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product to control any pests, by any application method or at any use sites, that are not specified on this label.

Use of this product may aid in the reduction of fleas, including by not necessarily limited to *Xenopsylla cheopis*, that commonly infest listed rodent hosts, if ingesting sufficient quantities of this bait.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, OR in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused or unconsumed bait as specified by this label.

RATS AND MICE

Use Restrictions: This product may be used to control Norway rats, roof rats and house mice at the sites in "Urban Areas" and "Non-urban Areas" that are indicated below. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes, and industrial, commercial and public buildings; in transport vehicles (ship, trains and aircraft) and in and around related port or terminal buildings. Applications "around" buildings must be made in tamper-resistant bait stations placed along walls or other external parts of buildings (e.g. doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

Non-urban Areas: To control Norway rats, roof rats and house mice this product may be used in and around homes and agricultural buildings. All outdoor placements must be in tamper-resistant bait stations.

Selection of Treatment Areas: Determine areas where rodents will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of rodents have been seen. Remove as much alternative food as possible.

Application Directions:

Norway Rats and Roof Rats: Apply 2 to 8 place packs per bait placement. Space bait placements at intervals of 15 to 30 feet in the infested area, at locations consistent with the requirements of this label. Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are fresh signs of rat activity or signs that rats are feeding on the bait.

House Mice: Open pack and apply 1/4 to 1/2 ounce of bait per typical placement. Larger placement – up to one unopened 2-ounce pack – may be made at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area, at locations consistent with the requirements on this label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

Follow-up (Rats & Mice): Immediately replace contaminated or spoiled bait. Collect and properly dispose of all visible dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

VOLES

Use Restrictions: This product may be used to control meadow voles (*Microtus pennsylvanicus*), pine voles (*M. pinetus*) and montane voles (*M. montanus*) around (within 15 feet of) homes, and industrial, commercial, agricultural and public buildings. Do not use this product within the habitat of threatened or endangered *Microtus* spp. voles, threatened or endangered *Peromyscus* spp. mice, or other threatened or endangered rodents. Do not expose bait on bare ground. Unless children, pets, birds and other wildlife are excluded from the treated areas, all applications for controlling voles must be made in tamper-resistant bait stations. All bait applications must be made by hand. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

Selection of Treatment Areas: Determine areas near buildings where voles or signs of vole activity (girdling of plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

Application Directions: Open pack and load one ounce of bait (half of a pack) into each bait

station. Situate bait stations near runways, burrow openings or plants that voles have been damaging. If, and only if, children, pets, domestic animals and wildlife reliably are excluded from the treated area, bait may be applied in one-ounce amounts directly into vole runways or burrow openings. Cover treated runways or holes with a shingle or board, or ground vegetation, if present. Check placements daily. Remove any fouled or spoiled bait, and replenish if vole activity and interest in bait persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of vole activity and feeding on bait. Collect and properly dispose of any carcasses and leftover bait.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Non-Refillable Container. Do not reuse or refill this container. Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration; or if allowed by state and local authorities, by burning; if burned, stay out of smoke.

WARRANTY

To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.

Scimetrics

LTD. CORP.

New Solutions to Old Problems

P.O. Box 1045 Wellington, CO 80549
(970) 482-1330

FOR CALIFORNIA: WARNING: THIS PRODUCT CONTAINS A CHEMICAL KNOWN TO THE STATE OF CALIFORNIA TO CAUSE BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM.

Label Version 4/2/08

